

NON COHEN ORACLE C.C.C.
SH669

SAHARON SHELAH

Institute of Mathematics
The Hebrew University
Jerusalem, Israel

Rutgers University
Mathematics Department
New Brunswick, NJ USA

ABSTRACT. The oracle c.c.c. is closely related to Cohen forcing. During an iteration we can “omit a type”; i.e. preserve “the intersection of a given family of Borel sets of reals is empty” provided that Cohen forcing satisfies it. We generalize this to other cases. In §1 we replace Cohen by “nicely” definable c.c.c., do the parallel of the oracle c.c.c. and end with a criterion for extracting a subforcing (not a complete subforcing, $\triangleleft$!) of a given nicely one and satisfying the oracle.

I would like to thank Alice Leonhardt for the beautiful typing.
First Typed - 1/9/98
Latest revision - 02/Nov/5

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\textsf{TeX}}$

§0 INTRODUCTION

This answers a question from [Sh:b, Ch.IV] (the chapter dealing with the oracle c.c.c.) asking to replace Cohen by e.g. random. Later we will deal with the parallel for oracle proper and for the case $\bar{\varphi}_\alpha$ is a (definition of a) nep forcing. An application will appear in a work with T. Bartoszynski.

How do we use this framework? We start with a universe satisfying $\diamond_{\aleph_1}$ and probably $2^{\aleph_1} = \aleph_2$ and choose $\langle S_i^* : i < \omega_2 \rangle$, $S_i^* \subseteq S^* \subseteq \omega_1$ such that $S_i^*/\mathcal{D}_{\omega_1}$ is strictly increasing and $\diamond_{S_{i+1}^* \setminus S_i^*}$ and for simplicity $S_i^* \subseteq S_{i+1}^*$ where $\mathcal{D}_{\omega_1}$ is the club filter on ω_1 . We choose by induction on $i < \omega_2$, a c.c.c. forcing $\mathbb{P}_i$ of cardinality $\aleph_1$ a sequence $\bar{M}^i = \langle M_\alpha^i : \alpha \in S_i^* \rangle$ and a 1-commitment.

They are increasing in the relevant sense and the work at limit stages is done by the general claims here. For $i = j + 1$ we have some freedom in choosing $\mathbb{P}_{i+1}$, usually $\mathbb{P}_{i+1} = \mathbb{P}_i * \underset{\sim}{\mathbb{Q}}_i$. So, working in $\mathbf{V}^{\mathbb{P}_i}$, $\mathbb{Q}_i$ has to satisfy a 0-commitment on S_i^* , and we like it to satisfy some task, possibly connected with some $X_i \subseteq \mathbb{R}^{V[\mathbb{P}_i]}$, say $X_i = X_i[G_{\mathbb{P}_i}]$ toward whatever task we have. We essentially have to choose $\bar{M}^i$ such that $\bar{M}^i \restriction S_j^* = \bar{M}^j$ but we have freedom to choose $\langle \bar{M}_\alpha^i : \alpha \in S_i^* \setminus S_j^* \rangle$ and a 0-commitment on $S_i^* \setminus S_j^*$. The reals generic for the chosen forcing notion as well as M_α^i for $\alpha \in S_i^* \setminus S_j^*$ can be chosen considering X_i . E.g. M_α^i can be the Mostowski Collapse of some $M \prec (\mathcal{H}(\aleph_2), \in)$ to which $\mathbb{P}_j, \bar{M}^j, x_j$ and X_j belongs.

So really this corresponds to the omitting type as in [Sh:e, XI].

§1 NON-COHEN ORACLE C.C.C.

1.1 Hypothesis.

(a) we assume CH, moreover $\diamond_{S^*}$ where $S^* \subseteq \{\delta < \omega_1 : \delta \text{ limit}\}$ is stationary.

1.2 Definition/Notation. 1) $\bar{M}$ denotes a sequence of the form $\langle M_\delta : \delta \in S \rangle$, M_δ a transitive countable model of ZFC^* , $\delta \subseteq M_\delta$, $M_\delta \models \text{"}\delta \text{ is countable"}$, $S \subseteq S^*$ stationary satisfying: for every $X \subseteq \omega_1$, the set $\{\delta \in S^* : X \cap \delta \in M_\delta\}$ is stationary. 2) $\mathcal{D}$ denotes a normal filter on ω_1 usually extending $\mathcal{D}_{\bar{M}}$ which is defined in 1.3(1) below (of course, the default value is $\mathcal{D}_{\bar{M}}$, see 1.4(1)).

We first give the old definitions from [Sh:f, IV]

1.3 Definition. 1) $\mathcal{D}_{\bar{M}}$ is

$\{X \subseteq \omega_1 : \text{for some } Y \subseteq \omega_1 \text{ we have : } \delta \in S_{\bar{M}} \cap X \Rightarrow Y \cap \delta \in M_\delta\}$.

2) A forcing notion $\mathbb{P}$ of cardinality $\leq \aleph_1$ satisfies the $(\bar{M}, \mathcal{D})$ -c.c. if for some (equivalently any) one to one $f : \mathbb{P} \rightarrow \omega_1$ the set:

$$\{\delta \in S_{\bar{M}} : \text{if } X \in M_\delta \text{ and } \{y \in \mathbb{P} : f(y) < \delta \text{ and } f(y) \in X\} \\ \text{is predense in } \mathbb{P} \upharpoonright \{y \in \mathbb{P} : f(y) < \delta\} \text{ then } X \text{ is predense in } \mathbb{P}\}$$

belongs to $\mathcal{D}$ and $\mathbb{P}$ has minimal element $\emptyset_{\mathbb{P}}$.

3) If $\mathcal{D} = \mathcal{D}_{\bar{M}}$ we may write " $\bar{M}$ -c.c.". Recall that $\mathcal{D}^+ = \{A \subseteq \omega_1 : \omega_1 \setminus A \notin \mathcal{D}\}$.

4) Let $\bar{M}^1 \leq \bar{M}^2$ if $\bar{M}^\ell = \langle M_\delta^\ell : \delta \in S_\ell \rangle$ and $\{\delta : \delta \in S_1 \setminus S_2 \text{ or } \delta \in S_1 \cap S_2 \text{ \& } M_\delta^1 \neq M_\delta^2\}$ is not stationary.

5) A forcing notion $\mathbb{P}$ satisfies the $(\bar{M}, \mathcal{D})$ -c.c. if: $|\mathbb{P}| \leq \aleph_0$ or for every $X \subseteq \mathbb{P}$ of cardinality $\leq \aleph_1$ there is $\mathbb{P}_1 \triangleleft \mathbb{P}$ of cardinality $\aleph_1$ which includes X and satisfies the $(\bar{M}, \mathcal{D})$ -c.c.

1.4 Fact. 1) $\mathcal{D}_{\bar{M}}$ is a normal filter on ω_1 .

2) The $\bar{M}$ -c.c. implies the c.c.c., and if $D_{\bar{M}} \subseteq \mathcal{D}$ (or just there is a normal filter $D' \supseteq D_{\bar{M}} \cup \mathcal{D}$) then the $(\bar{M}, \mathcal{D})$ -c.c.c. implies the c.c.c. and if $\mathcal{D}_2 \supseteq \mathcal{D}_1 \supseteq \mathcal{D}_{\bar{M}}$ are normal filters, then the $(\bar{M}, \mathcal{D}_1)$ -c.c. implies the $(\bar{M}, \mathcal{D}_2)$ -c.c.

3) We can find $\langle S_\zeta^* : \zeta < \omega_2 \rangle$ such that $S_\zeta^* \subseteq S^*$, $\zeta < \xi \Rightarrow S_\zeta^* \subseteq S_\xi^* \text{ mod } \mathcal{D}_{\bar{M}}$,

$S_\zeta^* \subseteq S_{\zeta+1}^*$ and $S_{\zeta+1}^* \setminus S_\zeta^* \in \mathcal{D}_{\bar{M}}^+$.

4) If $\bar{M}^1 \leq \bar{M}^2$ and the forcing notion $\mathbb{P}_2$ satisfies the $(\bar{M}^2, \mathcal{D})$ -c.c. and $\mathbb{P}_1 \triangleleft \mathbb{P}_2$, then $\mathbb{P}_1$ satisfies the $(\bar{M}^1, \mathcal{D})$ -c.c.

5) If $\bar{M}^1 \leq \bar{M}^2$ and $P_1 \triangleleft P_2$ and $\mathbb{P}_2$ satisfies the $\bar{M}^2$ -c.c. then $\mathbb{P}_1$ satisfies the $\bar{M}^1$ -c.c.

Proof. See [Sh:f, IV], but for the reader we prove one.

1) Without loss of generality $\mathbb{P}_2$ has set of elements ω_1 . As $\mathbb{P}_1 \triangleleft \mathbb{P}_2$ there is a function $f : \mathbb{P}_2 \rightarrow \mathbb{P}_1$ such that

(*)₁ $g \in \mathbb{P}_2 \wedge f(g) \leq_{\mathbb{P}_1} p \in \mathbb{P}_1 \Rightarrow p, g$ are compatible in $\mathbb{P}_2$.

Let $g : \mathbb{P}_2 \times \mathbb{P}_2 \rightarrow \mathbb{P}_2$ be such that

(*)₂ if $p, g \in \mathbb{P}_2$ are compatible then $g(p, g)$ is a common upper bound and $p, g \in \mathbb{P}_1 \Rightarrow g(p, g) \in \mathbb{P}_1$.

So there is a club E of ω_2 which is closed under f, g so

(*)₃ if $\delta \in S, \mathcal{J} \subseteq \mathbb{P}_1 \cap \delta$ is predense in $\mathbb{P}_1 \upharpoonright \delta$ then $\mathcal{J}$ is predense in $\mathbb{P}_2 \upharpoonright \delta$.
 [Why? If $g \in \mathbb{P}_2 \cap \delta$ then $f(g) \in \mathbb{P}_1 \cap \delta$ so by the assumption on $\mathcal{J}$, $f(g)$ is compatible with some $r_1 \in \mathcal{J}$ in $\mathbb{P}_1 \cap \delta$, so there is $r_2 \in \mathbb{P}_1 \upharpoonright \delta$ above g and r_1 . By the definition of f , r_2, g are compatible in $\mathbb{P}_2$ hence $g(r_2, g)$ is a common upper bound of them in $\mathbb{P}_2 \upharpoonright \delta$.]

1.5 Definition. 1) We say $\mathcal{V} = (S, \Phi, \bar{\eta}, \bar{\nu}) = (S^{\mathcal{V}}, \Phi^{\mathcal{V}}, \bar{\eta}^{\mathcal{V}}, \bar{\nu}^{\mathcal{V}})$ is a 0-commitment for $\bar{M}$ if for some $E \in \mathcal{D}_{\bar{M}}$:

- (a) $S \subseteq S^*, S \in \mathcal{D}_{\bar{M}}^+$
- (b) $\bar{\eta} = \langle \eta_\alpha : \alpha \in S \rangle, \Phi = \langle \bar{\varphi}_\alpha : \alpha \in S \rangle$ and if $\alpha \in S \cap E$ then $\bar{\varphi}_\alpha \in M_\alpha$ and $M_\alpha \models$ “ $\bar{\varphi}_\alpha$ is an absolute definition of a c.c.c. forcing notion called $\mathbb{Q}_\alpha$ with generic real η_α ”; note, absolute here means that forcing extensions of M_α , preserve predensity of countable sets (in the sense of M_α) order and incompatibility
- (c) $\bar{\nu} = \langle \nu_\alpha : \alpha \in S \rangle$ where $\nu_\alpha \in {}^\omega \omega$ and for every $\alpha \in S \cap E$ the real ν_α is $(\mathbb{Q}_\alpha, \eta_\alpha)$ -generic over M_α .

We ignore $\bar{M}$ if clear from the context. We can replace $\bar{M}$ by $(\bar{M}, \mathcal{D})$ if above $E \in \mathcal{D}, S \in \mathcal{D}^+$.

1A) A forcing notion $\mathbb{P}$ of cardinality $\leq \aleph_1$ satisfies the 0-commitment $\mathcal{V}$ if: $\mathbb{P}$ is an $\bar{M}$ -c.c. forcing notion and for any one-to-one mapping $h : \mathbb{P} \rightarrow \omega_1$ for some $E \in \mathcal{D}_{\bar{M}}$ we have

- (d) if $\alpha \in S \cap E$ then $\Vdash_{\mathbb{P}}$ “the real ν_α is a $(\mathbb{Q}_\alpha, \eta_\alpha)$ -generic real over $M_\alpha[\alpha \cap h''G_{\mathbb{P}}]$ ”.

2) Let $\mathbb{P} \in \mathcal{H}(\aleph_2)$ be an $\bar{M}$ -c.c. forcing notion. We say that $\mathcal{V} = (S, \bar{\Phi}, \bar{\eta}, \bar{\nu}) = (S^{\mathcal{V}}, \bar{\Phi}^{\mathcal{V}}, \bar{\eta}^{\mathcal{V}}, \bar{\nu})$ is a 1-commitment on $\mathbb{P}$ if: for any $\bar{N}$ satisfying $(*)_1$ below, the clauses (a)-(d) of $(*)_2$ below hold

- $(*)_1$ $\bar{N} = \langle N_\alpha : \alpha < \omega_1 \rangle$ is increasing continuous, $N_\alpha \prec (\mathcal{H}(\aleph_2), \in)$ is countable, $\bar{N} \upharpoonright (\alpha + 1) \in N_{\alpha+1}$ and $\{\bar{M}, \mathbb{P}\} \subseteq \bigcup_{\alpha < \omega_1} N_\alpha$
- $(*)_2$ (a) $S \subseteq \text{Dom}(\bar{M}) \subseteq S^*, S \in \mathcal{D}_{\bar{M}}^+$
- (b) $\bar{\eta} = \langle \eta_\alpha : \alpha \in S \rangle, \bar{\Phi} = \langle \bar{\varphi}_\alpha : \alpha \in S \rangle$ so $(\bar{\varphi}_\alpha, \eta_\alpha)$ is a $\mathbb{P}$ -name of a pair as in 1.5(1)(a), both are hereditarily countable over $\mathbb{P}$
- (c) $\bar{\nu} = \langle \nu_\alpha : \alpha \in S \rangle$ and ν_α a $\mathbb{P}$ -name of a real given by countably many conditions
- (d) the set of the $\alpha \in S$ satisfying the following belongs to $\mathcal{D}_{\bar{M}} + S$: $\bar{\varphi}_\alpha \in M_\alpha$, $\text{Mos Col}_{N_\alpha}(N_\alpha) \in M_\alpha$, and letting $\mathbb{P}'_\alpha = \text{Mos Col}_{N_\alpha}(N_\alpha^{\mathbb{P}}) = \{\text{Mos Col}_{N_\alpha}(x) : x \in \mathbb{P} \cap N_\alpha\} \in M_\alpha$ we have $M_\alpha \models \text{"}\bar{\varphi}_\alpha \text{ is a } \mathbb{P}'_\alpha\text{-name of an absolute definition of a c.c.c. forcing with generic real } \eta_\alpha\text{"}$ and $\Vdash_{\mathbb{P}} \text{"the real } \nu_\alpha \text{ is a } (\mathbb{Q}_{\bar{\varphi}_\alpha}, \eta_\alpha)\text{-generic real over } M_\alpha[G_{\mathbb{P}}]\text{"}$.

For simplicity the reader may concentrate on the case $\langle (\bar{\varphi}_\alpha, \eta_\alpha) : \alpha \in S \rangle \in \mathbf{V}$.

3) Let $IS = \{(\mathbb{P}, \mathcal{V}, \bar{M}) : \mathbb{P} \in \mathcal{H}(\aleph_2) \text{ is an } \bar{M}\text{-c.c. forcing notion and } \mathcal{V} \text{ is a 1-commitment on } \mathbb{P}\}$.

We shall omit $\bar{M}$ if clear from the context. We can replace $\bar{M}$ by $(\bar{M}, \mathcal{D})$ -naturally and write $IS_{\mathcal{D}}$, but the claims are the same.

4) For $(\mathbb{P}^\ell, \mathcal{V}^\ell, \bar{M}^\ell) \in IS$ ($\ell = 1, 2$) let $(\mathbb{P}^1, \mathcal{V}^1, \bar{M}^1) \leq^* (\mathbb{P}^2, \mathcal{V}^2, \bar{M}^2)$ means $\bar{M}^1 \leq \bar{M}^2$, $\mathbb{P}^1 \leq \mathbb{P}^2$ and for some $E \in \mathcal{D}_{\bar{M}^1}$ we have

$$S^{\mathcal{V}^1} \cap E \subseteq S^{\mathcal{V}^2} \cap E, \bar{\Phi}^{\mathcal{V}^1} \upharpoonright (S^{\mathcal{V}^1} \cap E) = \bar{\Phi}^{\mathcal{V}^2} \upharpoonright (S^{\mathcal{V}^2} \cap E),$$

$$\bar{\eta}^{\mathcal{V}^1} \upharpoonright (S^{\mathcal{V}^1} \cap E) = \bar{\eta}^{\mathcal{V}^2} \upharpoonright (S^{\mathcal{V}^2} \cap E) \text{ and } \bar{\nu}^{\mathcal{V}^1} \upharpoonright (S^{\mathcal{V}^1} \cap E) = \bar{\nu}^{\mathcal{V}^2} \upharpoonright (S^{\mathcal{V}^2} \cap E).$$

We call E a witness to $(\mathbb{P}^1, \mathcal{V}^1, \bar{M}^1) \leq^* (\mathbb{P}^2, \mathcal{V}^2, \bar{M}^2)$.

We point out the connection between 0-commitment and 1-commitment.

1.6 Fact: 1) If $\mathcal{V}$ is a 1-commitment on $\mathbb{P}$ and $\mathbb{P}$ an $\bar{M}$ -c.c. forcing notion of cardinality $\leq \aleph_1$, then $\Vdash_{\mathbb{P}} \text{"}\mathcal{V}[G_{\mathbb{P}}] = (S^{\mathcal{V}}, \bar{\Phi}^{\mathcal{V}}[G_{\mathbb{P}}], \bar{\eta}^{\mathcal{V}}[G_{\mathbb{P}}], \bar{\nu}[G_{\mathbb{P}}])\text{"}$ is a 0-commitment

so we call it $\mathcal{V}[G_{\mathbb{P}}]$. Note $\eta[G_{\mathbb{P}}]$ is still a name.

- 2) If $\mathbb{P} = \{\emptyset\}$ (the trivial forcing) then: $\mathcal{V}$ is a 1-commitment on $\mathbb{P}$ iff $\mathcal{V}$ is a 0-commitment.
- 3) If $\langle \bar{M}^i : i < \zeta \rangle$ is $\leq$ -increasing, $\zeta < \omega_2$ and $\text{Dom}(\bar{M}^i) \setminus S$ is not stationary, then there is $\bar{M}$, $\text{Dom}(\bar{M}) = S$ such that $i < \zeta \Rightarrow \bar{M}^i \leq \bar{M}$.
- 4) Increasing $\bar{M}$ preserves everything.

As a warm-up (see [Sh 630] for more)

1.7 Claim. 1) Assume

- (a) M is a countable transitive model of ZFC^- , $M \models \text{"}\mathbb{P}_1 \text{ is a countable forcing notion"}$
- (b) $M \models \text{"}\varphi \text{ is an absolute definition of c.c.c. forcing notion } \mathbb{Q}^\varphi \text{ with generic } \eta : \alpha_1 \rightarrow \alpha_2 \text{"}$
- (c) ν is $(M, \mathbb{Q}^\varphi)$ -generic sequence, i.e., there is $G \subseteq (\mathbb{Q}^\varphi)^M$ generic over M such that $\nu = \eta[G]$.

Then we can find a countable $\mathbb{P}_2$ such that

- (α) $\mathbb{P}_1 \subseteq_{ic} \mathbb{P}_2$ and every $\mathcal{J} \in M$ which is predense in $\mathbb{P}_1$ is predense in $\mathbb{P}_2$
- (β) $\Vdash_{\mathbb{P}_2} \text{"}\nu \text{ is } (M', \mathbb{Q}^\varphi)\text{-generic sequence where } M' = M[G_{\mathbb{P}_2} \cap \mathbb{P}_1]\text{"}$.

2) Similarly for φ defining a nep forcing.

Proof. 1) In M we can define $\mathbb{P}^+ = \mathbb{P}_1 * (\mathbb{Q}^\varphi)^{M[G_{\mathbb{P}_1}]}$, now as $\mathbb{Q}^\varphi$ is absolutely c.c.c., we know that $q \mapsto (\emptyset, q)$ is a complete embedding of $(\mathbb{Q}^\varphi)^M$ into $\mathbb{P}^+$. So if $G_* \subseteq (\mathbb{Q}^\varphi)^M$ is generic over M such that $\nu = \nu[G]$ then let $\mathbb{P}_2^* = \{(p, q) \in \mathbb{P}_1 * (\mathbb{Q}^\varphi)^{M[G_{\mathbb{P}_1}]} : (p, q) \text{ is compatible with } (\emptyset, q') \text{ for every } q' \in G_*\}$. Now check.
2) See [Sh 630]. $\square_{1.7}$

1.8 Crucial Claim. In IS , any $\leq^*$ -increasing ω -chain has an upper bound.

Proof. So assume $(\mathbb{P}^n, \mathcal{V}^n, \bar{M}^n) \in IS$ and $(\mathbb{P}^n, \mathcal{V}^n, \bar{M}^n) \leq^* (\mathbb{P}^{n+1}, \mathcal{V}^{n+1}, \bar{M}^{n+1})$ for $n < \omega$, let $\bar{M}$ be such that $\bar{M} \geq \bar{M}^n$ for each n ; so let $E_n \in \mathcal{D}_{\bar{M}}$ witness it. For simplicity assume that above any $p \in \mathbb{P}^n$ there are two incompatible elements, and $0 \in \mathbb{P}^0$ is minimal in all $\mathbb{P}^n$, i.e. is $\emptyset_{\mathbb{P}^n}$. Without loss of generality the set

of elements of $\mathbb{P}^n$ is $\subseteq \omega_1$ and $\omega_1 \setminus \bigcup_{n < \omega} \mathbb{P}^n$ has cardinality $\aleph_1$ and let X^* be such that $\bigcup_{n < \omega} \mathbb{P}^n \subseteq X^* \subseteq \omega_1$ and $|X^* \setminus \bigcup_{n < \omega} \mathbb{P}^n| = \aleph_1$; notation helps in a future use, also there we replace ω by a (countable) ordinal of cofinality $\aleph_0$. We can define functions $F_n, f_{n,m+1}, F_{n,m,\ell}$ (when $n < m < \omega, \ell < \omega$) such that

- (a)_n if $p, q \in \mathbb{P}_n$ are compatible then $F_n(p, q) \in \mathbb{P}_n$ is a common upper bound
- (b)_{n,m} if $n \leq m$ and $p \in \mathbb{P}_m$, then $\langle F_{n,m,\ell}(p) : \ell < \omega \rangle$ is a maximal antichain of $\mathbb{P}_n$, such that for each ℓ :
either $p, F_{n,m,\ell}(p)$ are incompatible (in $\mathbb{P}_m$)
or p is compatible in $\mathbb{P}_m$ with every $q \in \mathbb{P}_n$ which is above $F_{n,m,\ell}(p)$
- (c)_{n,m} if $p \in \mathbb{P}_m, g \in \mathbb{P}_n$ then $F_{n,m}(p, g) \in \mathbb{P}_n$ and if there is $r, g \leq_{\mathbb{P}_n} r$ and r, p is incompatible in $\mathbb{P}_m$ then $p, F_{n,m}(p, g)$ are incompatible in $\mathbb{P}_m$.

Let E be a club of ω_1 , such that $\delta \in E \Rightarrow \delta$ is closed under $F_n, F_{n,m,\ell}$ and $\text{otp}(X^* \cap \delta \setminus \bigcup_{n < \omega} \mathbb{P}^n) = \delta$.

We would like to define a forcing notion $\mathbb{P}^\omega$ with universe X^* , and 1-commitment $\mathcal{Y}^\omega$, and functions $F_\omega, F_{n,\omega,\ell}$ satisfying the natural requirements. First, let $E_\omega = \bigcap_{n < \omega} E_n \cap E, S^{\mathcal{Y}^\omega} = \bigcup_{n < \omega} S^{\mathcal{Y}^n} \cap E_\omega$, and for $\alpha \in S^{\mathcal{Y}^\omega}$ the triple $(\tilde{\varphi}_\alpha^{\mathcal{Y}^\omega}, \tilde{\eta}_\alpha^{\mathcal{Y}^\omega}, \tilde{\nu}_\alpha^{\mathcal{Y}^\omega})$ is $(\tilde{\varphi}_\alpha^{\mathcal{Y}^{n(\alpha)}}, \tilde{\eta}_\alpha^{\mathcal{Y}^{n(\alpha)}}, \tilde{\nu}_\alpha^{\mathcal{Y}^{n(\alpha)}})$ where $n(\alpha) = \text{Min}\{n : \alpha \in S^{\mathcal{Y}^n}\}$.

Defining $\mathbb{P}_\omega, F_\omega, F_{n,\omega,\ell}$ is harder, so we first define AP , a set of approximations to it. A member t of AP has the form $(\delta^t, \mathbb{P}^t, F_\omega^t, F_{n,\omega,\ell}^t, \Gamma^t)_{\ell < \omega}$ satisfying

- (α) $\delta^t \in E$
- (β) $\mathbb{P}^t$ is a forcing notion with set of elements $\subseteq X^* \cap \delta^t$ and $\supseteq \delta^t \cap \bigcup_n \mathbb{P}_n$ and $0 \leq_{\mathbb{P}^t} p$ for every $p \in \mathbb{P}^t$
- (γ) $\mathbb{P}^t \restriction (\mathbb{P}^n \cap \delta^t) = \mathbb{P}^n \restriction (\mathbb{P}^n \cap \delta^t)$
- (δ) if $p, q \in \mathbb{P}^t$ are compatible in $\mathbb{P}^t$ then $F_\omega^t(p, q)$ is such an upper bound
- (ε) if $p \in \mathbb{P}^t, n < \omega$ then $\langle F_{n,\omega,\ell}^t(p) : \ell < \omega \rangle$ is a maximal antichain of $\mathbb{P}_n$, the members are $< \delta^t$, and for each ℓ , either $p, F_{n,\omega,\ell}^t(p)$ are incompatible in $\mathbb{P}^t$ or $(\forall q \in \mathbb{P}_n \cap \delta)(\mathbb{P}_n \models "F_{n,\omega,\ell}(p) \leq q" \Rightarrow p, q \text{ are compatible in } \mathbb{P}^t)$ and for at least one ℓ the second case occurs
- (ζ) if $p \in \mathbb{P}^t \cap \mathbb{P}_m \setminus \bigcup_{\ell < m} \mathbb{P}_\ell$ then $F_{n,\omega,\ell}^t(p) = F_{n,m,\ell}(p)$

- (η) Γ^t is a sequence $\langle \bar{p}_\zeta^t : \zeta < \zeta^t \rangle$, $\zeta^t < \omega_1$ and $\bar{p}_\zeta^t$ is a sequence of length $\leq \omega$ of members of $\mathbb{P}^t$ which form a maximal antichain (of $\mathbb{P}^t$)
- (θ) if $p \in \mathbb{P}^t$ and $n < m < \omega$ and $r \in \mathbb{P}_n \cap \delta^t$ and $[r \leq r' \in \mathbb{P}_n \cap \delta^t \Rightarrow r', p \text{ are compatible in } \mathbb{P}^t]$, then the set $\{F_{m,\omega,\ell}^t(p) : \ell < \omega \text{ and } p \text{ is compatible with } F_{m,\omega,\ell}^t(p) \text{ in } \mathbb{P}^t\}$ satisfies: if $r \leq g \in \mathbb{P}_n$ then in $\mathbb{P}_m$, g is compatible with some member of this set
- (ι) if $\zeta < \zeta^t$ and $n < \omega$ then:
 $\{F_{n,\omega,\ell}^t(p_{\zeta,k}^t) : k < \omega, \ell < \omega \text{ and } p_{\zeta,k}^t, F_{n,\omega,\ell}^t(p_{\zeta,k}^t) \text{ are compatible in } \mathbb{P}^t\}$ is a predense subset of $\mathbb{P}_n$. Note that trivial, this subset is predense in $\mathbb{P}_n \cap \delta$; similarly in clause (κ)

Moreover,

- (κ) if $p^* \in \mathbb{P}^t$ and $n < \omega$ and $\zeta < \zeta^t$ then

$$\begin{aligned} \mathcal{J}_{\zeta,n,p^*}^t =: \{r' \in \mathbb{P}_n \cap \delta : & (i) \quad r', p^* \text{ incompatible in } \mathbb{P}^t \text{ or} \\ & (ii) \quad \text{for some } k < \omega \text{ and } p' \text{ we have} \\ & (\forall r'')[r' \leq r'' \in \mathbb{P}_n \cap \delta \rightarrow \{r'', p'\} \\ & \text{has an upper bound in } \mathbb{P}^t] \text{ and } p^* \leq_{\mathbb{P}^t} p', p_{\zeta,k}^t \leq_{\mathbb{P}^t} p'\}. \end{aligned}$$

is predense in $\mathbb{P}_n$.

* * *

We define the (natural) partial order $\leq^*$ on AP : for $t, s \in AP$ as follows; we let $t \leq^* s$ iff:

- (i) $\delta^t \leq \delta^s$
- (ii) $\mathbb{P}^t \subseteq \mathbb{P}^s$
- (iii) $F_\omega^t \subseteq F_\omega^s$
- (iv) $F_{n,\omega,\ell}^t \subseteq F_{n,\omega,\ell}^s$
- (v) Γ^t is an initial segment of Γ^s .

Fact A: $AP \neq \emptyset$.

Proof. Easy: choose $\delta \in E$, let $\mathbb{P}^t = (\bigcup_{n < \omega} \mathbb{P}^n) \restriction \delta$, $F_\omega(p, q) = F_{n(p,q)}(p, q)$ where

$$n(p, q) = \text{Min}\{n : p \in \mathbb{P}_n \text{ and } q \in \mathbb{P}_n\}.$$

For $n < \omega, p \in \mathbb{P}^t \cap \delta$ let $\langle F_{n,\omega,\ell}^t(p) : \ell < \omega \rangle$ be $\langle F_{n,m,\ell}(p) : \ell < \omega \rangle$ for the first

$m \geq n$ such that $p \in \mathbb{P}_m$.

Lastly, $\Gamma =$ empty sequence.

Fact B: If $t \in AP$ and $\delta^t < \delta \in E$, then there is s satisfying $t \leq^* s \in AP$ with $\delta^s \geq \delta, \zeta^s = \zeta^t$.

Proof. Without loss of generality $t, \langle \mathbb{P}^n \restriction \delta : n < \omega \rangle, X^* \cap \delta$ belongs to M_δ and $\delta \in E \cap \bigcap_{n < \omega} E_n$ and $X^* \cap \delta \setminus \bigcup_{n < \omega} \mathbb{P}_n \setminus \delta^t$ is infinite. [Why? As $\emptyset \notin \mathcal{D}_{\bar{M}}$ and we can increase δ .] So (for the last phrase see the proof of 1.4(4))

(*) any $\mathcal{J} \in M_\delta$ which is a predense subset of $\mathbb{P}_n \restriction \delta$ is a predense subset of $\mathbb{P}_n$ and $n < m \Rightarrow \mathbb{P}_n \restriction \delta \leq \mathbb{P}_m \restriction \delta$.

Let $A = \mathbb{P}^t \subseteq X^* \cap \delta^t, B = \bigcup_{n < \omega} \mathbb{P}_n \cap \delta$. We define a forcing notion $\mathbb{Q}$, with set of elements $\subseteq A \times B$ identifying $(p, 0)$ with p and $(0, q)$ with q . Now $(p, q) \in A \times B$ belongs to $\mathbb{Q}$ iff: $p = 0$ or $q = 0$ or there are $r \in A \cap B$ and $n = n(p, q)$ such that: $\mathbb{P}_n \models "r \leq q"$, and $(\forall r')[r \leq r' \in \mathbb{P}_n \cap \delta^t \rightarrow r', p \text{ compatible in } \mathbb{P}^t]$; we call such r a witness and n a possible value for $n(p, q)$. The order on $\mathbb{Q}$ is $(p, q) \leq (p', q') \Leftrightarrow p \leq_{\mathbb{P}^t} p' \ \& \ \bigvee_n q \leq_{\mathbb{P}_n} q'$.

Now note

(α) $\mathbb{Q} \restriction A = \mathbb{P}^t$

(β) $\mathbb{Q} \restriction B = \bigcup_{n < \omega} \mathbb{P}_n \restriction \delta$

(γ) If $(p, q) \in \mathbb{Q}, m = n(p, q)$ and $q \in \mathbb{P}_m \restriction \delta$ and $\mathbb{P}_m \restriction \delta \models "q \leq q'"$ and $\mathbb{P}^t \models "p' \leq p"$, then $(p', q') \in \mathbb{Q}$ and $\mathbb{Q} \models "(p, q) \leq (p', q')"$

(δ) if $(p, q) \in \mathbb{Q}$ and $n = n(p, q) \leq m < \omega$, then for some q_1 we have: $(p, q) \leq (p, q_1) \in \mathbb{Q}$ and $n(p, q_1) = m$, or at least m is a possible value for $n(p, q_1)$. [Why? Let $r \in \mathbb{P}_{n(p, q)}$ be a witness in particular r is compatible with p in $\mathbb{P}^t$. By clause (θ) of the Definition of AP the set $\mathcal{J} = \{F_{m, \omega, \ell}^t(p) : \ell < \omega \text{ and } p \text{ is compatible with } F_{m, \omega, \ell}^t(p) \text{ in } \mathbb{P}^t\}$ is predense above r in $\mathbb{P}_m$. $\mathbb{P}_n \models r \leq g$ hence $\mathbb{P}_m \models r \leq g$ so for some $\ell, F_{m, \omega, \ell}^t(p) \in \mathcal{J}$ is compatible with q in $\mathbb{P}_m$ so there is $q_1 \in \mathbb{P}_m \cap \delta$ such that $\mathbb{P}_m \models q \leq q_1 \wedge F_{m, \omega, \ell}^t(p) \leq q_1$. So $(p, q_1) \in \mathbb{Q}$ as witnessed by m and $r' = F_{m, \omega, \ell}^t(p)$, is as required.]

(ε) $\mathbb{P}_n \restriction \delta \leq \mathbb{Q}$

[Why? Let $(p^0, q^0) \in \mathbb{Q}$, of course, we can replace this pair by any larger one, so by clause (δ) above without loss of generality some $m \in [n, \omega)$, is a

possible value for $n(p^0, q^0)$ so we have $q^0 \in \mathbb{P}_m \restriction \delta$, hence recalling that $\mathbb{P}_n \restriction \delta \leq \mathbb{P}_m \restriction \delta$ there is $q^1 \in \mathbb{P}_n \restriction \delta$ such that:

$$(\forall r \in \mathbb{P}_n)(\mathbb{P}_n \restriction \delta \models q^1 \leq r \Rightarrow r, q^0 \text{ compatible in } \mathbb{P}_m \restriction \delta).$$

Assume $q^1 \leq r \in \mathbb{P}_n \restriction \delta$. So r, q^0 are compatible in $\mathbb{P}_m \restriction \delta$ hence has a common upper bound $q^2 \in \mathbb{P}_m \restriction \delta$.

In particular $q^0 \leq q^2 \in \mathbb{P}_m \restriction \delta$ so by clause (γ) we have $(p^0, q^2) \in \mathbb{Q}$ and $(p^0, q^0) \leq^Q (p^0, q^2)$; also $r = (0, r) \leq (p^0, q^2)$ as $r \leq q^2$ together $r, (p^0, q^0)$ are compatible in $\mathbb{Q}$, so $[q^1 \leq r \in \mathbb{P}_n \restriction \delta \Rightarrow (p^0, q^0), r = (0, r) \text{ are compatible in } \mathbb{Q}]$. As $(p^0, q^0) \in \mathbb{Q}$ was arbitrary we are done.]

(ζ) if $p_1, p_2 \in \mathbb{P}^t$ are incompatible in $\mathbb{P}^t$ then they are incompatible in $\mathbb{Q}$.
[Why? Look at the order of $\mathbb{Q}$.]

(η) if $\zeta < \zeta^t$ then $\bar{p}_\zeta^t$ is a maximal antichain in $\mathbb{Q}$.

[Why? If not some $(p^*, q^*) \in \mathbb{Q}$ is incompatible in $\mathbb{Q}$ with every $(p_{\zeta, k}^t, 0)$ for $k < \omega$. Let $n < \omega$ be a possible value of $n(p^*, q^*)$ so $q^* \in \mathbb{P}_n \restriction \delta$ and there is a witness $r^* \leq q^*, r^* \in \mathbb{P}_n \restriction \delta^t$ for $(p^*, q^*) \in \mathbb{Q}$.

By clause (κ) in the definition of $t \in AP$ we know that for some $r \in \mathbb{P}_n \cap \delta^t$ we have:

$$(i) \quad r \in \mathcal{I}_{\zeta, n, p^*}^t$$

$$(ii) \quad q^*, r \text{ are compatible in } \mathbb{P}_n$$

As q^*, r are compatible and $r^* \leq q^*$ also r^*, r are compatible in $\mathbb{P}_n$ hence in $\mathbb{P}_n \cap \delta^t$, so by the demand on r^* , we have: r, p^* are compatible in $\mathbb{P}^t$. So in clause (ι) of definition of AP , in the definition of $\mathcal{I}_{\zeta, n, p^*}^t$ for our r subclause (i) fails hence subclause (ii) holds so there are k, p' as in subclause (ii) there. Also let $q^1 \in \mathbb{P}_n \restriction \delta$ be a common upper bound of q^*, r . So r witness that $(p', q^1) \in \mathbb{Q}$ with n a possible value of $n(p', q^1)$. Clearly it is above (p^*, q^*) and above $p_{\zeta, k}^t$ so we are done.]

Let $\delta^s = \delta$. Clearly $\mathbb{Q} \in M_\delta$ and recall that $M_\delta \models \text{"}\delta \text{ is countable, } \mathbb{P}^t \text{ is countable"}$ so there is $f \in M_\delta$ such that $f : \mathbb{Q} \rightarrow X^* \cap \delta$ is a one to one (into or even onto), extending $\text{id}_A \cup \text{id}_B$, and define $\mathbb{P}^s$ such that f is an isomorphism from $\mathbb{Q}$ onto $\mathbb{P}^s$. We can define $F_\omega^s, F_{n, \omega, \ell}^s (n, \ell < \omega)$ extending $F_\omega^t, F_{n, \omega, \ell}^s$ as required, e.g., $F_{n, \omega, \ell}^s((p, q)) = F_{n, m, \ell}^s(g)$ for some $m > n$ such that $g \in \mathbb{P}_m$ except when $g = 0$ then $F_{n, \omega, \ell}^s((p, 0)) = F_{n, \omega, \ell}^t(p)$. Now it is easy to check clause (θ) of the definition of $s \in AP$, recalling $(*)$ above and clauses $(i), (\kappa)$ holds since the construction is made in M_δ . Lastly, let $\Gamma^s = \Gamma^t$.

Fact C: If $t^n \in AP$ and $t^n \leq^* t^{n+1}$ for $n < \omega$ then there is t such that $n < \omega \Rightarrow t^n \leq^* t \in AP$ and $\delta^t = \bigcup_{n < \omega} \delta^{t^n}$ and $\zeta^t = \bigcup_{n < \omega} \zeta^{t^n}$.

[Why? Just let δ^t, ζ^t be as above, $\mathbb{P}^t = \bigcup_{n < \omega} \mathbb{P}^{t^n}$, $F_\omega^t = \bigcup_{n < \omega} F_n^{t^n}$, $F_{m,\omega,\ell}^t = \bigcup_{n < \omega} F_{m,\omega,\ell}^{t^n}$ and $p_{\zeta,k}^t = p_{\zeta,k}^{t^n}$ for every n large enough. Now check.]

Main Fact D: Assume $t \in AP$, $\delta^t \in E \cap \bigcap_{n < \omega} E_n$, $t \in M_\delta$ and $\delta^t \in \bigcup_{n < \omega} S^{\mathcal{Y}_n}$. Then there is $s \in AP$ such that $t \leq^* s$ and ν_δ is actually a $\mathbb{P}^s$ -name (i.e. all the countably many conditions appearing in its definition belongs to $\bigcup_{n < \omega} \mathbb{P}_n \cap \delta^t \subseteq \mathbb{P}^t$) and:

- (*) if $\mathbb{P}^t \subseteq_{ic} \mathbb{Q}$, and for each $\zeta < \zeta^s$ the sequence $\bar{p}_\zeta^t$ is a maximal antichain of $\mathbb{Q}$, then
 $\Vdash_{\mathbb{Q}}$ “there is $G' \subseteq \mathbb{Q}_{\bar{\varphi}_\delta^t}^{M_\delta[G]}$ generic over $M_\delta[G]$ such that $(\eta[G])(G') = \nu_\delta$ ”.

[Why? Chase arrows.]

So we can choose $t_\varepsilon \in AP$ by induction on $\varepsilon < \omega_1$ such that t^ε is $\leq^*$ -increasing continuous, $\delta^{t^{\varepsilon+1}} > \delta^{t^\varepsilon}$, and if $t^\varepsilon \in M_{(\delta^{t^\varepsilon})}$, $\delta^{t^\varepsilon} \in \bigcap_{n < \omega} E_n \cap E \cap \bigcup_{n < \omega} S^{\mathcal{Y}_n}$ then $t^{\varepsilon+1}$ is gotten by Fact D. No problem to carry this ($\varepsilon = 0$ by Fact A, $\varepsilon = \varepsilon_1 + 1$ by Fact D if possible and by Fact B if not; lastly, if ε is a limit ordinal, use Fact C).

Now let $\mathbb{P}^\omega = \bigcup_{\varepsilon < \omega_1} \mathbb{P}^{t^\varepsilon}$ and it should be clear how to define $\mathcal{Y}^\omega$; now check the requirements. $\square_{1.8}$

1.9 Definition. Let $\bar{C}^* = \langle C_\delta^* : \delta < \omega_2 \text{ a limit ordinal} \rangle$ (and $C_\alpha^* = \emptyset$ otherwise) be a square and $\bar{X}^* = \langle X_i^* : i < \omega_1 \rangle$ be an increasing sequence of subsets of ω_1 , $|X_i^* \setminus \bigcup_{j < i} X_j^*| = \aleph_1$, $X_{\omega_1}^* = \bigcup_{i < \omega_1} X_i^*$.

We say that $\langle (\mathbb{P}_i, \mathcal{Y}_i, f_i, \bar{M}_i) : i < \alpha \rangle$ is a $(\bar{C}^*, \bar{X}^*)$ -iteration (we omit $\bar{M}^i$ and write $(\bar{M}, \bar{C}^*, \bar{X}^*)$ -iteration if $i < \alpha \Rightarrow \bar{M}^i \leq \bar{M}$ or an $\bar{M}$ -iteration when $\bar{C}^*, \bar{X}$ are clear from context) if:

- (a) $(\mathbb{P}_i, \mathcal{Y}_i, \bar{M}^i) \in IS$ is $<^*$ -increasing and $\text{Dom}(\bar{M}^i) = S^{\mathcal{Y}_i}$
- (b) f_i is a one to one function from $\mathbb{P}_i$ onto $X_{\text{otp}(C_\alpha^*)}^*$, and let $(\mathbb{P}'_i, \mathcal{Y}'_i)$ be such that f_i maps $(\mathbb{P}_i, \mathcal{Y}_i)$ to $(\mathbb{P}'_i, \mathcal{Y}'_i)$
- (c) if $j \in \text{acc}(C_i)$ then $f_j \subseteq f_i$
- (d) if $\text{cf}(i) = \aleph_0$ and $i = \sup \text{acc}(C_i^*)$ then $(\mathbb{P}'_i, \mathcal{Y}'_i)$ is gotten from $\langle (\mathbb{P}'_j, \mathcal{Y}'_j) : j \in \text{acc}(C_i^*) \rangle$ as in the proof of 1.8 (using $\langle X_j^* : j \in \text{acc}(C_i^*) \rangle, X_{\text{otp}(C_i^*)}^*$)

- instead of $\langle X_n : n < \omega \rangle, X_\omega$ so $\text{acc}(C_i^*)$ replace ω and we generate $\langle t_\alpha^i : \alpha < \omega_1 \rangle$ and by it define $(\mathbb{P}'_i, \mathcal{Y}'_i)$ hence $(\mathbb{P}_i, \mathcal{Y}_i)$
- (e) in clause (d), assume $\delta = \text{otp}(C_i^*), \langle \langle \mathbb{P}'_j, \mathcal{Y}'_j \rangle \restriction \delta : j \in \text{acc}(C_i^*) \rangle \in M_\delta$ and for $j_1 < j_2$ from $\text{acc}(C_i^*)$ the ordinal δ belongs to the club $\{\alpha < \omega_1 : \alpha \text{ limit closed under the functions } F^{j_1} \text{ and } F_\ell^{j_1, j_2} \text{ (see clause (f) below)}\}$ and $\delta^{t_\delta^{j_1}} = \delta$. Let $t_*^i \in AP$ be defined by $\delta_*^i = \text{otp}(C_i^*), \mathbb{P}^{t_*^i} = \cup \{\mathbb{P}'_j \restriction \delta : j \in \text{acc}(C_i^*)\}, F_{\omega^*}^{t_*^i} = \cup \{F^{j_1} \restriction \delta_{i, j_1} \in \text{acc}(C_i^*)\}, F_{j_1, \omega, \ell}^{t_*^i} = \cup \{F_\ell^{j_1, j_2} : j_2 \in \text{acc}(C_i^*) \setminus j_1\}$ and let $\Gamma^{t_*^i}$ be empty. If $t_*^i \in M_\delta$ then let t_0^i be gotten from t_*^i as in Fact (D).
- (f) F^j is a (partial) two-place function from $X_{\text{otp}(C_j^*)}^*$ to itself such that $F^j(p, q)$ is the $<$ -first common upper bound of p and q in $\mathbb{P}'_j$ and if $j_1 \in \text{acc}(C_{j_2}^*)$ then we have $\langle F_n^{j_1, j_2}(p) : n < \omega \rangle$ is a maximal antichain of $\mathbb{P}'_{j_1}$ satisfying: for each n , either $F_n^{j_1, j_2}(p)$ is incompatible with p in $\mathbb{P}'_{j_2}$ or p is compatible with r in $\mathbb{P}'_{j_2}$ wherever $\mathbb{P}'_{j_1} \models F_n^{j_1, j_2}(p) \leq r$.

1.10 Claim. (iteration at limit) 1) Assume $\langle (\mathbb{P}_i, \mathcal{Y}_i, f_i) : i < \zeta \rangle$ is a $(\bar{M}, \bar{C}^*, \bar{X}^*)$ -iteration where $\zeta < \omega_2$ is a limit ordinal. Then

- (a) we can find $(\mathbb{P}_\zeta, \mathcal{Y}_\zeta, f_\zeta)$ such that $\langle (\mathbb{P}_i, \mathcal{Y}_i, f_i) : i < \zeta + 1 \rangle$ is an $\bar{M}$ -iteration
- (b) if $S \subseteq S^*, i < \zeta \Rightarrow S^{\mathcal{Y}_i} \subseteq S \text{ mod } \mathcal{D}_{\bar{M}}, \text{ then we can demand } S^{\mathcal{Y}_\zeta} = S$.

Proof. If $\text{cf}(\zeta) = \aleph_0$ we use 1.8 but taking care of clause (e), this just dictates to us how to start the induction there. If $\text{cf}(\zeta) = \aleph_1$, then by the square bookkeeping (see clause (e) in Definition 1.9) our work is done (using $f_\zeta = \cup \{f_\xi : \xi \in \text{acc}(C_\zeta)\}$). $\square_{1.10}$

1.11 Claim. 1) Assume

- (a) $\mathcal{Y} = (S, \bar{\Phi}, \bar{\eta}, \bar{\nu})$ is a 1-commitment on the forcing notion $\mathbb{P} \in \mathcal{H}(\aleph_2)$ for $\bar{M}$
- (b) $G_\mathbb{P} \subseteq \mathbb{P}$ is generic over $\mathbf{V}, \bar{\nu}^0 = \langle \nu_\alpha^0 : \alpha \in S \rangle$ where $\nu_\alpha^0 = \nu_\alpha[G_\mathbb{P}]$, $\bar{M}^1 = \bar{M}[G_\mathbb{P}] = \langle M_\delta[f'', G_\mathbb{P}] : \delta \in S^* \rangle$ for some one to one function f from $\mathbb{P}$ into ω_1
- (c) in $\mathbf{V}[G_\mathbb{P}]$, $\mathcal{Y}^1 = (S^1, \bar{\Phi}^1, \bar{\eta}^1, \bar{\nu}^1)$ is a 0-commitment, $S \subseteq S^1 \text{ mod } \mathcal{D}_{\bar{M}[G_\mathbb{P}]}, \bar{\Phi}^1 \restriction (S \cap S^1) = \bar{\Phi} \restriction (S \cap S_1)$,

$$\begin{aligned} \bar{\eta}^1 \restriction (S \cap S_1) &= \bar{\eta} \restriction (S \cap S_1), \bar{\nu}^1 \restriction (S \cap S_1) = \bar{\nu}^0 \restriction (S \cap S_1) \text{ and} \\ (\bar{S}^1, \bar{\Phi}^1, \bar{\eta}^1) &\in V \end{aligned}$$

(d) in $\mathbf{V}[G_{\mathbb{P}}]$, $\mathbb{Q}$ is a forcing notion satisfying the 0-commitment $\mathcal{Y}^1$ for $\bar{M}^1$.

Then for some $\mathbb{P}$ -name $\mathbb{Q}$ and 1-commitment $\mathcal{Y}^2$ we have:

- (a) $(\mathbb{P}, \mathcal{Y}) \leq^* (\mathbb{P} * \mathbb{Q}, \mathcal{Y}^2)$
- (b) $S^{\mathcal{Y}^2} = S^1, \Phi^{\mathcal{Y}^2} = \bar{\Phi}^1, \bar{\eta}^{\mathcal{Y}^2} = \bar{\eta}^1, \bar{\nu}[G_P] = \bar{\nu}^1$
- (c) $\mathbb{Q}[G_{\mathbb{P}}] = \mathbb{Q}$.

2) If for every $G_{\mathbb{P}} \subseteq \mathbb{P}$ generic over $\mathbf{V}$ there are $\mathbb{Q}$ satisfying some ψ_1 and $(S^1, \bar{\Phi}^1, \bar{\eta}^1, \bar{\nu}) \in \mathbf{V}[G_{\mathbb{P}}]$ as above satisfying some ψ_2 , then we can demand

(d) $\Vdash_{\mathbb{P}} \text{“}\mathbb{Q}[G_{\mathbb{P}}], \mathcal{Y}^2 \text{ as above satisfies } \psi_1, \psi_2 \text{ respectively”}$.

3) We may allow $\langle (\bar{\varphi}_{\alpha}, \bar{\eta}_{\alpha}) : \alpha \in S^1 \rangle$ be a sequence of $\mathbb{P}$ -names and even $(\mathbb{P} * \mathbb{Q})$ -names.

Proof. Straight.

1.12 Claim. (iteration in successor case: increase the commitment).

Assume $\langle (\mathbb{P}_i, \mathcal{Y}_i, f_i) : i < \zeta \rangle$ is an $\bar{M}$ -iteration and $\zeta = \xi + 1, S^{\mathcal{Y}_{\xi}} \subseteq S \subseteq S^*, S \subseteq \text{Dom}(\bar{M})$ and $\langle (\bar{\varphi}_{\alpha}, \bar{\eta}_{\alpha}) : \alpha \in S \setminus S^{\mathcal{Y}_{\xi}} \rangle$ is as required in Definition 1.5.

Lastly $Z_{\alpha} \subseteq {}^{\omega}2$ is a $\mathbb{P}_{\xi}$ -name of a positive set for $(\bar{\varphi}_{\alpha}, \bar{\eta}_{\alpha})$ for every such α .

Then we can find $(\mathbb{P}_{\zeta}, \mathcal{Y}_{\zeta}, f_{\zeta})$ such that

- (i) $\langle (\mathbb{P}_i, \mathcal{Y}_i, f_i) : i < \zeta + 1 \rangle$ is an $\bar{M}$ -iteration
- (ii) $\mathbb{P}_{\zeta} = \mathbb{P}_{\xi}, S^{\mathcal{Y}_{\zeta}} = S, (\bar{\varphi}_{\alpha}^{\mathcal{Y}_{\zeta}}, \bar{\eta}_{\alpha}^{\mathcal{Y}_{\zeta}}) = (\bar{\varphi}_{\alpha}, \bar{\eta}_{\alpha})$ if $\alpha \in S \setminus S^{\mathcal{Y}_{\xi}}$.

Proof. Straight.

1.13 Claim. (*iteration at successor: increasing the forcing*)

Suppose

- (a) $(\mathbb{P}, \mathcal{Y}) \in IS$ and the set of elements of $\mathbb{P}$ is X_i (the X_j^* 's as in 1.9)
- (b) $\mathbb{Q}$ is a $\mathbb{P}$ -name satisfying, for every $G \subseteq \mathbb{P}$ generic over $\mathbf{V}$, the following:
 - (i) $\mathbb{Q}[G]$ is a forcing notion with set of elements beings $X_{i+1} \setminus X_i$
 - (ii) $\{\delta < \omega_1 : \text{if } \mathbb{P} \restriction \delta \in M_\delta \text{ and } G \cap \delta \text{ is a generic subset of } \mathbb{P} \restriction \delta, \mathbb{Q}[G] \restriction \delta \in M_\delta[G \cap \delta] \text{ and } \nu_\delta[G] \text{ is forced to be generic for } ((\mathbb{Q}^{\varphi_\delta[G]})^{M_\delta[G]}, \eta_\delta[G])\} \in \mathcal{D}_{\bar{M}[G]}$

Then we can find $(\mathbb{P}^+, \mathcal{Y}^+)$ such that $(\mathbb{P}, \mathcal{Y}) \leq^* (\mathbb{P}^+, \mathcal{Y}^+) \in IS$ and the $\mathbb{P}$ -name $\mathbb{P}^+/G_{\mathbb{P}}$ is equivalent to $\mathbb{Q}[G_{\mathbb{P}}]$.

Proof. Straight.

1.14 Conclusion. Assume $(\bar{C}^*, \bar{X}^*)$ is as in 1.8. Let Φ be a set of definitions of forcing notions with some real parameters, and $\langle S_i^* : i < \omega_2 \rangle$ is as in 1.4 for $\mathcal{D}_{\bar{M}}$.

We can find $\langle (\mathbb{P}_i, \mathcal{Y}_i, f_i, \bar{M}^i) : i < \omega_2 \rangle$ such that

- (a) it is an $(\bar{C}^*, \bar{X}^*)$ -iteration
- (b) $\mathbb{P} = \bigcup_{i < \omega_2} \mathbb{P}_i$ is a c.c.c. forcing notion of cardinality $\aleph_2$ (so in $\mathbf{V}^{\mathbb{P}}, 2^{\aleph_0} \leq \aleph_2$) and except in degenerated cases equality holds)
- (c) $S^{\mathcal{Y}_i} = S_i^*$ from 1.4(3)
- (d) if in $\mathbf{V}^{\mathbb{P}_i}$ we have $(\bar{\varphi}, \eta)$ is a case of Φ as in 1.5, moreover
 - $\Vdash_{\mathbb{P}_i} \{ \delta \in S_{i+1}^* \setminus S_i^* : M_\delta^{i+1}[f_i'', G_{\mathbb{P}_i}] \models "(\bar{\varphi}, \eta) \text{ as required in 1.5}" \} \in \mathcal{D}_{\bar{M}^{i+1}}^+$ (even less with more bookkeeping) and $Z \subseteq (\omega_2)^{\mathbf{V}^{\mathbb{P}}}$ is positive for $(\bar{\varphi}, \eta)$, then
 - (α) $\{ \delta \in S^{\mathcal{Y}_{i+1}} \setminus S^{\mathcal{Y}_i} : (\bar{\varphi}, \eta_\delta)/G_{\mathbb{P}_i} = (\bar{\varphi}, \eta) \text{ and } \nu_\delta[G_{\mathbb{P}_i}] \in Z \} \in \mathcal{D}_{\bar{M}}^+$, in fact the set is forced to include such old set (from $\mathbf{V}$) by this we can get
 - (β) for some $j > i, \delta \in S^{\mathcal{Y}_{j+1}} \setminus S_j^{\mathcal{Y}_0} \Rightarrow (\bar{\varphi}, \eta_\delta)/G_{\mathbb{P}_i} = (\bar{\varphi}, \eta), \nu_\delta[G_{\mathbb{P}_i}] \in Z$

- (e) if H is a pregiven function such that for every $i < \omega_2$ and $(\mathbb{P}, \mathcal{Y}, \bar{M})$ satisfying
- $(\mathbb{P}_i, \mathcal{Y}_i) \leq^* (\mathbb{P}, \mathcal{Y}) \in IS$ such that $S^{\mathcal{Y}} = \mathcal{S}_i$ we have $(\mathbb{P}, \mathcal{Y}) \leq^* H(\mathbb{P}, \mathcal{Y}) \in IS$ such that $H(\mathbb{P}, \mathcal{Y}, \bar{M})$ satisfies the demands from (a) + (c) on $(\mathbb{P}_{i+1}, \mathcal{Y}_{i+1}, \bar{M}^{i+1})$, then we can demand $(\exists^{\aleph_2} j)[(\mathbb{P}_{j+1}, \mathcal{Y}_{j+1}) = H(\mathbb{P}_j, \mathcal{Y}_j)]$; moreover, if $S^* \subseteq \omega_2$ is stationary we can demand $\{j \in S : (\mathbb{P}_{j+1}, \mathcal{Y}_{j+1}) = H(\mathbb{P}_j, \mathcal{Y}_j)\}$ is stationary.
- (Of course, we can promise this for $\aleph_2$ such functions).

Proof. Put together the previous claims. (Concerning clause (e) without loss of generality $\{i < \omega_1 : \text{otp}(C_i^*) = 0\}$ is stationary) so in those stages we have no influence of clause (e) of 1.9; anyhow the influence of 1.9(e) is minor.

1.15 Discussion. We discuss here some possible extensions.

1.16 Claim. Assume $\langle S_i : i < \omega_2 \rangle$ is a sequence of pairwise almost disjoint stationary subsets of ω_1 , each with diamond and $i < j \Rightarrow S_i \subseteq S_j^+ \bmod \mathcal{D}_{\omega_1}$, so $S_i^+ \subseteq \omega_1$ and $S_i \cap S_i^+ = \emptyset$.

Then in the following game the between the bookkeeper and the forcer, the bookkeeper has a winning strategy.

A Play last ω_2 moves, before the α -th move a sequence $\langle (\mathbb{P}_i, \mathbb{Q}_i, \bar{M}_i, \mathcal{Y}) : i < \alpha \rangle$ is defined such that

- (a) $\mathbb{P}_i$ a c.c.c. forcing notion of cardinality $\aleph_1$, say $\subseteq \mathcal{H}_{<\aleph_1}(\aleph_2)$
- (b) $\mathbb{Q}_i$ is a $\mathbb{P}_i$ -name of a forcing notion of cardinality $\leq \aleph_1$, say $\subseteq \omega_1$
- (c) $\mathbb{P}_i$ is $\prec$ -increasing
- (d) $\mathbb{P}_{i+1}, \mathbb{P}_i * \mathbb{Q}_i$ are isomorphic over $\mathbb{P}_i$
- (e) $\bar{M}_i$ is a $\mathbb{P}_i$ -name of an S_i -oracle
- (f) $\mathcal{Y}_i$ is a $\mathbb{P}_i$ -name of a S_i -commitment.

In the i -th move:

the bookkeeper chooses $\mathbb{P}_i$ and a $\mathbb{P}_i$ -name $(\bar{M}_i^+, \mathcal{Y}_i^+)$ of an S_i^+ -oracle and 0-commitment the forces choose $\mathbb{Q}_i$ and $(\bar{M}_i, \mathcal{Y}_i)$, $\mathbb{P}_i$ -names such that $\mathbb{Q}_i$ satisfies $(\bar{M}_i^+, \mathcal{Y}_i^+)$ and $(\bar{M}_i, \mathcal{Y}_i)$.

In the end the bookkeeper wins if

$$i < j < \omega_2 \Rightarrow \mathbb{P}_j/\mathbb{P}_i \text{ satisfies } (\bar{M}_i, \mathcal{Y}_i).$$

Proof. Similar to earlier proofs.

We give an easy criterion for existence.
The following uses more from [Sh 630].

1.17 Claim. *Assume*

- (a) $(\mathbb{P}, \leq, \leq_n)_{n < \omega}$ is a definition of a forcing notion satisfying condition A of Baumgartner with $\leq_n$ as witness and ZFC_*^- says this, in a way preserved by suitable forcing
- (b) $\mathcal{Y}$ is a 0-commitment,
- (c) $\mathbb{P}$ is absolutely nep such that for each $\alpha \in S^{\mathcal{Y}}$ it is $\leq_n$ -purely $I_{\mathbb{Q}^{\bar{\varphi}\alpha}}$ -preserving, i.e.
 - (*) if M is a $\mathbb{P}$ -candidate and a $\mathbb{Q}^{\bar{\varphi}\alpha}$ -candidate, $p \in \mathbb{P}^M$, $n < \omega$ and $q \in (\mathbb{Q}^{\bar{\varphi}\alpha})^M$ then for some p', η, ν we have $p \leq_n p' \in \mathbb{P}$, p' is $\langle M, \mathbb{P} \rangle$ -generic and ν is $(\mathbb{Q}^{\bar{\varphi}\alpha}, \eta_\alpha)$ -generic over M satisfying q (check def) and $p' \Vdash_{\mathbb{P}} \text{“}\nu \text{ is } (\mathbb{Q}^{\bar{\varphi}\alpha}, \eta_\alpha)\text{-generic over } M \langle G_{\mathbb{P}} \cap P^M \rangle\text{”}$.

Then there is a c.c.c. forcing notion $\mathbb{P}' \subseteq \mathbb{P}$ (not necessarily $\mathbb{P}' \triangleleft \mathbb{P}$) satisfying the 0-commitment $\mathcal{Y}$ and $\Vdash_{\mathbb{P}} \text{“for a club of } \delta < \omega_1, \varphi(\nu, \eta_\delta^*)\text{”}$.

Remark. 1) Why the φ_δ 's? We hope it helps, for example in the following; suppose we are given $f : \mathbb{R} \rightarrow \mathbb{R}$, we like to force $A \subseteq \mathbb{R}$ which is not in $I_{\mathbb{Q}^{\bar{\varphi}\alpha}}$ and on which the function f is continuous; i.e. to force a continuous f^* such that $\{\eta \in {}^\omega 2 : f^*(\eta) = f(\eta)\} \in (I_{\mathbb{Q}}^{ex})^+$. So not only do we like to find $q \Vdash \text{“}\eta_\delta \text{ is } (\mathbb{Q}_\delta, \eta_\delta)\text{-generic over } M_\delta[G_{\mathbb{P}}]\text{”}$ but also $q \Vdash_{\mathbb{P}'} \text{“}f(\eta_\delta) = f(\eta_\delta^*)\text{”}$. This is what $\bar{\varphi}$ says.

Proof. We choose by induction on $\alpha < \omega_1$, a pair $(\mathbb{P}_\alpha, \Gamma_\alpha)$ such that:

- (α) $\mathbb{P}_\alpha \subseteq \mathbb{P}$ is countable
- (β) Γ_α is a countable family of predense subsets of $\mathbb{P}_\alpha$

- (γ) if $\mathcal{J} \in \Gamma_\alpha$ and $p \in \mathbb{P}_\alpha$ and $n < \omega$ then for some q we have $p \leq_n q \in \mathbb{P}_\alpha$ and $\mathcal{J}$ is predense above q in $\mathbb{P}$
- (δ) $\mathbb{P}_\alpha$ is increasing continuous in α
- (ε) Γ_α is increasing continuous in α .

Case 1: $\alpha = 0$. Trivial.

Case 2: $\alpha = \beta + 1$, β non-limit or $(\mathbb{P}_\beta, \Gamma_\beta) \notin M_\beta$.

Let $(\mathbb{P}_\alpha, \Gamma_\alpha) = (\mathbb{P}_\beta, \Gamma_\beta)$.

Case 3: α limit.

Let $(\mathbb{P}_\alpha, \Gamma_\alpha) = (\bigcup_{\beta < \alpha} \mathbb{P}_\beta, \bigcup_{\beta < \alpha} \Gamma_\beta)$.

Case 4: $\alpha = \delta + 1$ where δ is a limit ordinal and $(\mathbb{P}_\delta, \Gamma_\delta) \in M_\delta$.

We can find $g \subseteq \text{Levy}(\aleph_0, |\mathbb{P}|)^{M_\delta}$, generic over M_δ such that η_δ^* is still $\mathbb{Q}_\delta$ -generic over $M_\delta[g]$ (see [Sh 630], §6).

In $M_\delta[g]$ we define $\mathbb{P}_\delta^+ = \{p : M_\delta[g] \models p \in \mathbb{P} \text{ and } \mathcal{J} \in \Gamma_\alpha \Rightarrow \mathcal{J} \text{ predense above } p'\}$ using the induction hypothesis, as in $M_\delta[g]$ the sets Γ_δ is countable:

(*) for every $p \in \mathbb{P}_\delta$ and $n < \omega$ there is $p' \in \mathbb{P}_\delta^+$ such that $\mathbb{P} \models p \leq_n p'$.

Again by [Sh 630], §6 for every $n < \omega$ and $p \in \mathbb{P}_\delta^+$, there is $q_{p,n} \in \mathbb{P}$ such that $p \leq_n q_{p,n} \in \mathbb{P}$, $q_{p,n}$ is $(M_\delta[g], \mathbb{Q})$ -generic and $q_{p,n} \Vdash_{\mathbb{P}} \text{"}\eta_\delta \text{ is a } (\mathbb{Q}_\delta, \eta_\delta)\text{-generic real over } M_\delta[g][G_{\mathbb{P}}] \text{ and } \varphi_\delta(\nu, \eta_\delta)\text{"}$.

Let $\mathbb{P}_{\delta+1} = \mathbb{P}_\delta \cup \{q_{p,n} : p \in \mathbb{P}_\delta^+ \text{ and } n < \omega\}$ and $\Gamma_{\delta+1} = \Gamma_\delta \cup \{\mathcal{J}_\delta\}$ where $\mathcal{J}_\delta = \{q_{p,n} : p \in \mathbb{P}_\delta^+ \text{ and } n < \omega\}$.

□_{1.17}

REFERENCES.

- [Sh:e] Saharon Shelah. *Non-structure theory*, accepted. Oxford University Press.
- [Sh 630] Saharon Shelah. Non-elementary proper forcing notions. *Journal of Applied Analysis*, **accepted**. math.LO/9712283.
- [Sh:b] Saharon Shelah. *Proper forcing*, volume 940 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, xxix+496 pp, 1982.
- [Sh:f] Saharon Shelah. *Proper and improper forcing*. Perspectives in Mathematical Logic. Springer, 1998.